



Theoretical exploration of amentoflavone and bilobetin interactions with BPIFB4: implications for longevity and cardioprotection

Ivan Vito Ferrari^{1*} 

¹Institute of Clinical Physiology, National Research Council, Via Aurelia Sud, 54100 Massa, Italy. E-mail: ivanvitoferrari@gmail.com

Article Info

Volume 1, Issue 1, January 2024

Received : 13 September 2023

Accepted : 29 December 2023

Published: 25 January 2024

doi: [10.62587/AFRJBS.1.1.2024.53-58](https://doi.org/10.62587/AFRJBS.1.1.2024.53-58)

Abstract

This study delves into the therapeutic potential of two natural compounds, amentoflavone and bilobetin, by investigating their interactions with the BPIFB4 protein. BPIFB4, a member of the BPI fold-containing superfamily, has been associated with various biological processes, particularly in longevity and cardioprotection (LAV-BPIFB4). Molecular docking analyses revealed promising binding energies for amentoflavone and bilobetin with BPIFB4, suggesting favorable interactions. Additionally, by pKCSM database are showed the predicted toxicity properties of amentoflavone and bilobetin respectively. This research contributes to the growing body of knowledge on natural compounds and their therapeutic implications in the context of vital biological processes. However, these findings underscore the need for further investigations, including in vivo studies and clinical trials, to validate the observed effects and explore the translational potential of amentoflavone and bilobetin in promoting cardiovascular health and longevity.

Key words: LAV-BPIFB4, Amentoflavone, Bilobetin, Docking analysis, pKCSM

© 2024 Ivan Vito Ferrari. This is an open access article under the CC BY license (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

1. Introduction

BPI fold-containing family B member 4 (BPIFB4) is a human protein encoded by the BPIFB4 gene, formerly identified as “palate, lung, and nasal epithelial carcinoma-associated protein 4” from the LPLUNC4 gene. The BPIFB4 gene produces four transcripts (splice variants), with three well-characterized isoforms. This protein is prominently expressed in the olfactory epithelium (nasal mucosa) and exhibits elevated levels in the gonads (testis, ovary) and the pituitary gland. Additionally, moderate levels are detected in white blood cells, particularly monocytes. BPIFB4 can be either localized within the cell cytoplasm or secreted, circulating systemically in the blood plasma. These findings suggest diverse functions for BPIFB4 in various tissues, emphasizing significant expression in olfactory and gonadal tissues and the potential for circulation in the bloodstream (Villa *et al.*, 2018; Bingle *et al.*, 2011; Ciaglia *et al.*, 2020). BPIFB4 belongs to the BPI fold protein superfamily, characterized by the presence of the bactericidal/permeability-increasing protein fold (BPI fold). This fold is created by two similar domains arranged in a “boomerang” shape. The BPI fold is a structural motif often found in proteins involved in host defense mechanisms and innate immune responses, and it plays a role in interactions with bacteria and other pathogens. The “boomerang” shape refers to the characteristic structure formed by these domains, contributing to the functional properties of proteins within this superfamily (Beamer *et al.*, 1998).

* Corresponding author: Ivan Vito Ferrari, Institute of Clinical Physiology, National Research Council, Via Aurelia Sud, 54100 Massa, Italy. E-mail: ivanvitoferrari@gmail.com

Analyzing numerous centenarian populations has enabled the identification of the longevity-associated variant (LAV) within the BPIFB4 gene. Individuals exhibiting elevated circulating levels of LAV-BPIFB4 seem to experience cardioprotective effects and display immunomodulatory properties, particularly in terms of anti-inflammatory responses (Dossena *et al.*, 2020; Cattaneo *et al.*, 2023; Montella *et al.*, 2021).

This study represents the inaugural exploration of this protein using Molecular Docking (Fan *et al.*, 2019) using Autodock Vina (Trott and Olson, 2010) with Pyrx program[10], aiming to identify the most effective molecule for enhanced binding.


A preliminary Virtual Screening study using the Pyrx program (Dallakyan and Olson, 2015) scrutinized hundreds of drugs and natural molecules. This process identified potential lead compounds with the capability to interact with this protein.

2. Materials and methods

BPIFB4 protein was performed by Autodock Vina (Trott and Olson, 2010) with Pyrx program (Dallakyan and Olson, 2015). This protein was taken by AlphaFold from Uniprot Database (BPIFB4 - BPI fold-containing family B member 4 - Homo sapiens (Human) | UniProtKB | UniProt). Given the unavailability of this protein in the Protein Data Bank, the AlphaFoldDB-predicted version was employed.

The docking study focused solely on portions of the protein with high Model Confidence (pLDDT > 90) and areas with very good Model Confidence (90 > pLDDT > 70). Protein loop areas with low Model Confidence were omitted and excluded from consideration in the Molecular Docking analysis (See below Figure 1).

- BPI fold containing family B, member 4 (AlphaFold),BPIFB4_HUMAN); Grid box Coordinates of Binding Center X (-4.0306), Y(0.8353), Z(0.3699); Size_x = 114.84863636 Size_y = 47.8852510452size_Z= 77.6464200592 to perform Blind Docking analysis.



3. Results and Discussion

The BPIFB4 protein is typically present in the olfactory epithelium, mononuclear cells, macrophage-like cells, and various progenitor/stem cell types. In the olfactory mucosa, it is thought to function similarly to other BPIFB proteins by contributing to the innate immune response when exposed to bacteria. When circulating in

the bloodstream, BPIFB4 is presumed to engage with endothelial cells that line blood vessels, leading to vasorelaxation in the smooth muscle cells of blood vessels (Dossena *et al.*, 2020). The examination of various groups of centenarians has led to the discovery of the longevity-associated variant (LAV) within the BPIFB4 gene. Those with heightened circulating levels of LAV-BPIFB4 demonstrate apparent cardioprotective effects and exhibit immunomodulatory features, specifically with anti-inflammatory characteristics (Dossena *et al.*, 2020; Cattaneo *et al.*, 2023; Montella *et al.*, 2021).

In this groundbreaking research, two potential candidates were identified through Docking analysis (Fan *et al.*, 2019; Trott and Olson, 2010), demonstrating a high binding energy score (approximately -9.5 kcal/mol) for interaction with BPIFB4. This discovery follows comprehensive preliminary Virtual Screening studies (Dallakyan and Olson, 2015) that assessed hundreds of drugs and natural molecules.

The outcomes of the docking analysis are presented in Figures 3 and 4, elucidating the particular chemical bonds formed by amentoflavone, a biflavonoid, and bilobetin, a flavonoid oligomer, in interaction with the BPIFB4 protein.

The noteworthy molecules identified in this study are two natural substances: amentoflavone and bilobetin. They are two highly similar molecules from a chemical structure perspective.

Figure 2 illustrates their comparison.




Figure 2: Comparison 3D structure of amentoflavone with 3D structure bilobetin respectively




Figure 3: Displays the docking outcomes of BPI fold-containing family B member 4 in conjunction with amentoflavone within as analyzed by Autodock Vina through Pyrx program. On the left side, 2D diagrams illustrate the residue interactions between the protein and amentoflavone. Meanwhile, the right side exhibits the potential Ligand Binding Site of the protein, highlighting the specific location of amentoflavone




Figure 4: Displays the docking outcomes of BPI fold-containing family B member 4 in conjunction with bilobetin within, as analyzed by Autodock Vina through Pyrx program. On the left side, 2D diagrams illustrate the residue interactions between the protein and bilobetin. Meanwhile, the right side exhibits the potential Ligand Binding Site of the protein, highlighting the specific location of bilobetin

Table 1: Shows the comparison of predicted toxicity parameters with amentoflavone and bilobetin through pKCSM Server

Compounds	AME Stoxicity	Max. tolerated dose(human) (log mg/kg/day)	hERG I inhibitor	hERG II inhibitor	Oral Rat Acute Toxicity (LD50) (mol/kg)	Oral Rat Chronic Toxicity (LOAEL) (log mg/kg_bw/day)	Hepatotoxicity
Amentoflavone	no	0.438	no	yes	2.527	3.572	no
Bilobetin	no	0.437	no	yes	2.56	2.217	no

In summary, the docking analysis revealed that bilobetin, a distinct flavonoid present in *Ginkgo biloba* leaves, exhibited a binding energy of approximately -9.4 kcal/mol with the BPIFB protein. Conversely, amentoflavone, characterized as a biflavonoid consisting of two linked flavonoid units, demonstrated a binding energy of approximately -9.5 kcal/mol with the BPIFB protein.

The observed characteristics of both bilobetin and amentoflavone, along with their interactions with the crucial BPIFB4 protein, position them as promising candidates for biological applications. This is especially noteworthy considering the involvement of BPIFB4 in various biological processes, particularly in relation to the longevity-associated variant (LAV) of the BPIFB4 gene.

The identification of LAV through the study of centenarian populations suggests that individuals with elevated circulating levels of LAV-BPIFB4 may experience cardioprotective effects (Sultana et al., 2022; Dewey et al., 2023) and an immunodulatory response, characterized by anti-inflammatory properties (Cattaneo et al., 2023; Montella et al., 2021).

It could be interesting conducting in-depth studies on substances like amentoflavone and bilobetin is crucial to understanding their therapeutic effects and potential activation or stimulation of BPIFB4 protein. Some aspects that researchers may explore in such studies:

3.1. Mechanism of interaction

Investigate the molecular mechanisms by which amentoflavone and bilobetin interact with BPIFB4.

3.2. Functional effects

Examine the downstream effects of BPIFB4 activation or stimulation induced by amentoflavone and bilobetin.

Assess the impact on cellular and physiological processes related to longevity and cardioprotectiveness.

3.3. Therapeutic potential

Evaluate the therapeutic potential of amentoflavone and bilobetin in conditions related to cardiovascular health, considering BPIFB4's suggested cardioprotective effects.

Explore their potential in anti-inflammatory and immunomodulatory contexts.

3.4. Bioavailability and pharmacokinetics

Investigate the bioavailability and pharmacokinetics of amentoflavone and bilobetin to understand how efficiently these substances can reach and interact with BPIFB4 in different tissues.

3.5. Cellular and animal studies

Conduct cellular and animal studies to validate the findings obtained from molecular docking studies.

3.6. Clinical trials

If preclinical studies are promising, consider progressing to clinical trials to evaluate the safety and efficacy of amentoflavone and bilobetin in humans, particularly in the context of longevity and cardiovascular health.

Given the potential implications for health and longevity, continued research on these substances and their interactions with BPIFB4 could contribute valuable insights to both basic science and clinical applications.

4. Conclusion

This research delves into the therapeutic potential of natural compounds amentoflavone and bilobetin, investigating their interactions with the BPIFB4 protein—a pivotal member of the BPI fold-containing superfamily associated with diverse biological processes, particularly longevity and cardioprotection (LAV-BPIFB4). Molecular docking analyses indicate promising binding energies for amentoflavone and bilobetin with BPIFB4, suggesting favorable interactions. Moreover, the pKCSM database sheds light on the predicted toxicity properties of these compounds, contributing to their understanding of their safety profiles. In pKCSM toxicity predictions, amentoflavone is more favorable than bilobetin with an Oral Rat Acute Toxicity (LD50) of 2.527 mol/kg vs. 2.56 mol/kg. Both exhibit comparable values for Max. Tolerated Dose (human) and Oral Rat Chronic Toxicity (LOAEL), indicating similar overall toxicity profiles. Nevertheless, it underscores the need for further investigations, including in vivo studies and clinical trials, to validate observed effects and explore the translational potential of amentoflavone and bilobetin in enhancing cardiovascular health and promoting longevity. The comprehensive exploration of these compounds holds promise for advancing this understanding of their safety and efficacy, potentially paving the way for new therapeutic avenues.

References

- Beamer, L.J., Carroll, S.F. and Eisenberg, D. (1998). [The BPI/LBP family of proteins: a structural analysis of conserved regions. *Protein Science*, 7\(4\): 906-914.](#)
- Bingle, C.D., Seal, R.L. and Craven, C.J. (2011). [Systematic nomenclature for the PLUNC/PSP/BSP30/SMGB proteins as a subfamily of the BPI fold-containing superfamily. *Biochem Soc Trans*, 39\(4\): 977-983.](#)
- Dossena, M., Ferrario, A., Lopardo, V., Ciaglia, E. and Puca, A.A. (2020). [New insights for BPIFB4 in cardiovascular therapy. *International Journal of Molecular Sciences*, 21\(19\); 7163.](#)

- Cattaneo, M., Beltrami, A.P., Thomas, A.C., Spinetti, G., Alvino, V.V., Avolio, E. and Madeddu, P. (2023). The longevity-associated BPIFB4 gene supports cardiac function and vascularization in ageing cardiomyopathy. *Cardiovasc Res.*, 119(7): 1583-1595. doi: 10.1093/cvr/cvad008.
- Ciaglia, E., Montella, F., Lopardo, V., Scala, P., Ferrario, A., Cattaneo, M. and Puca, A.A. (2020). Circulating BPIFB4 levels associate with and influence the abundance of reparative monocytes and macrophages in long living individuals. *Frontiers in Immunology*, 11; 1034.
- Dallakyan, S. and Olson, A.J. (2015). Small-molecule library screening by docking with PyRx. *Chemical Biology: Methods and Protocols*, 243-250.
- Fan, J., Fu, A. and Zhang, L. (2019). Progress in molecular docking. *Quantitative Biology*, 7; 83-89.
- Montella, F., Lopardo, V., Cattaneo, M., Carrizzo, A., Vecchione, C., Ciaglia, E. and Puca, A.A. (2021). The role of BPIFB4 in immune system and cardiovascular disease: The lesson from centenarians. *Translational Medicine@ UniSa*, 24(1); 1.
- Trott, O. and Olson, A. J. (2010). AutoDock Vina: improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading. *Journal of computational chemistry*, 31(2); 455- 461.
- Villa, F., Carrizzo, A., Ferrario, A., Maciag, A., Cattaneo, M., Spinelli, C.C. and Puca, A. A. (2018). A model of evolutionary selection: The cardiovascular protective function of the longevity associated variant of BPIFB4. *International Journal of Molecular Sciences*, 19(10); 3229.

Cite this article as: Ivan Vito Ferrari (2024). Theoretical exploration of amentoflavone and bilobetin interactions with bpihb4: implications for longevity and cardioprotection. *African Research Journal of Biosciences*. 1(1), 53-58. doi: 10.62587/AFRJBS.1.1.2024.53-58.